

Treating Pigs for Round Worm

RECOMMENDATIONS

Use sodium fluoride mixed with damp or dry mash or ground grain for single or group treatment of pigs infested with round worm.

DOSAGE

Live weight of Pigs	Sodium Fluoride	Number of Pigs
20 lb.	1 oz. treats	18
21-40 lb.	1 oz. "	9
41-60 lb.	2 oz. "	10
61-80 lb.	2 oz. "	8
81-100 lb.	3 oz. "	9
101-150 lb.	3 oz. "	7
151-200 lb.	5 oz. "	9
Over 200 lb.	5 oz. "	7

In the event of Poisoning send for a Doctor.

FIRST AID TREATMENT

Give abundant drinks of water or milk.

Give lime water or chalk.

Give stimulants.

Apply artificial respiration.

Wash eyes and skin with a stream of water for 15-30 minutes.

WARNING

Sodium Fluoride if used at the above dosage is safe. Excessive dosage may lead to poor bone and tooth structure. Do not dose a pig more than four or five times during its life.

POISON S5

KEEP OUT OF REACH OF CHILDREN



SODIUM FLUORIDE

FOR ROUND WORM IN PIGS

Active Constituent:

98% SODIUM FLUORIDE

Avoid contact of the skin and eyes.

NET CONTENTS: / LB.

Imported and packed by

A.C.F. & Shirleys Fertilizers Ltd.

Little Roma Street, Brisbane

DIRECTIONS FOR USE

Weigh the animals to determine the exact dosage.

Starve on the evening before treatment. Allow pigs plenty of water.

Carefully weigh out correct dosage and thoroughly mix with damp or dry mash or well ground grain.

Allow 1 lb. dry weight of food for each 25 lb. live weight of pig.

Feed treated food in morning and leave until all has been consumed.

Choose food with high lime content but do not use milk, garbage or whole grain.

WARNING

The use of Sodium Fluoride being beyond the control of A.C.F. and Shirleys Fertilizers Ltd., no warranty expressed, implied or inferred, is made with respect to the suitability or fitness of that product for purposes for which it is to be used by the Buyer, whether or not in accordance with any directions furnished or claimed so to be, and the Buyer assumes responsibility for any results whatsoever which might occur.

R.P. 1M. 3/50.